



Orthodontic Treatment Information Sheet

Orthodontic Treatment Orthodontic treatment promotes a healthy smile, healthier teeth and gums and a better bite for normal chewing. Orthodontic treatment depends for much of its success on the understanding and co-operation of the patient. This form covers the usual material risks associated with orthodontic treatment. Other factors may also be described by your orthodontist. You may ask your orthodontist to explain anything you do not understand. While recognising the benefits of healthy teeth and a pleasing smile, you should also be aware that orthodontic treatment has limitations and potential risks and there may be alternative treatment options. You should clarify what is expected of you as a patient, or as the parent of a young patient, to achieve excellent results. Keep in mind that with orthodontic treatment, like other healing arts, results cannot be guaranteed. The unknown factor in any orthodontic correction is the response of the patient to the orthodontic treatment.

Should you have any questions, please ask.

Benefits Orthodontics plays an important role in improving overall oral health and achieving balance and harmony between the face and teeth for a healthy smile, which may enhance one's self-esteem. The correct bite of the teeth with healthy muscles and jaw joints will provide optimal chewing function. Properly aligned teeth are easier to brush and so the tendency to decay may be decreased, as may the likelihood of developing disease of the gums and supporting bone. Correctly aligned teeth are less likely to experience abnormal wear. Because of the individual conditions present and the limitations of treatment imposed by nature each specific benefit may not be attainable for every patient.

Risks All forms of medical and dental treatment including orthodontics have some risks and limitations. Fortunately in orthodontics complications are infrequent and when they do occur they are usually of only minor consequence. Nevertheless, they should be considered when making the decision to undergo orthodontic treatment. The principal risks in orthodontic treatment pertain to:

1. **Oral Hygiene.** Orthodontic appliances do not cause tooth decay. Gum disease, tooth decay and permanent markings (decalcification) on the teeth can occur if orthodontic patients eat food containing excessive sugar and high acid drinks or do not brush their teeth frequently or properly. These same problems can occur without orthodontic treatment but the risk is greater to the individual wearing braces. It is imperative during orthodontic treatment that extra care be taken with oral hygiene and tooth brushing in particular.
2. **Relapse.** Teeth may have a tendency to change their positions after treatment. The long-term, faithful wearing of retainers should reduce this tendency. Teeth can however, move at any time, whether or not they had orthodontic treatment. This is especially true during the late teen period when active growth of the facial bones is coming to an end. The most vulnerable teeth are those at the front. Throughout life the bite can change adversely from various causes such as eruption of wisdom teeth, growth or maturational changes, mouth breathing, playing on musical instruments and other oral habits. Very occasionally tooth movements will be severe enough to merit a further course of treatment to ensure a satisfactory result.
3. **Periodontal Health.** The health of the bone and gums which support the teeth may be affected by orthodontic tooth movement if the condition already exists, and in some rare cases where a condition does not appear to exist. In general, orthodontic treatment lessens the possibility of tooth loss or gum infection due to misalignment of the teeth or jaws. Inflammation of the gums and loss of supporting bone can occur if bacterial plaque is not removed daily with good oral hygiene.
4. **Root shortening.** In some patients the length of the roots of the teeth may be shortened during orthodontic treatment. Some patients are prone to this happening, some are not. It is nearly impossible to predict susceptibility. Usually the shortening does not have significant consequences, but on rare occasions it may become a threat to the longevity of the teeth involved.
5. **Jaw Joints.** Occasionally, patients may suffer pain or dysfunction in the jaw joints (TMJ). This may present as joint pain, headaches or ear problems. These problems may occur with or without orthodontic treatment. Any of the above noted symptoms should be reported to the orthodontist. Some patients are susceptible to TMJ problems and some are not. Numerous studies have shown that the movement of teeth during orthodontic treatment is unrelated to the onset of TMJ problems. Just as for painful conditions in other joints, discomfort in TMJ can last from a few days to several weeks or longer.
6. **Tooth vitality.** Sometimes a tooth may have been traumatised by a previous accident or a tooth may be decayed or have fillings which can cause damage to the nerve of the tooth. Orthodontic tooth movement in some cases aggravates this condition and in rare instances may lead to loss of tooth vitality and discolouration of the tooth requiring root canal treatment and other dental treatment to restore the colour of the tooth.

7. **Loose appliances and discomfort.** The gums, cheeks or lips may be scratched or irritated by newly placed appliances or by loose or broken appliances or by blows to the mouth. You will be given instructions on minimising these effects. Very rarely loose orthodontic appliances may be accidentally swallowed or aspirated. You should inform your orthodontist of any unusual symptoms, or broken or loose appliances, as soon as they are noted. Usually post adjustment tenderness should be expected and the period of tenderness or sensitivity varies with each patient and the procedure performed. (Typical post-adjustment tenderness may last 24 to 48 hours).
8. **Atypical growth.** Insufficient excessive or asymmetrical changes in the growth of the jaws may limit the orthodontist's ability to achieve the desired result. If growth becomes disproportionate during orthodontic management the treatment may be prolonged. Growth changes that occur after orthodontic treatment may alter the quality of treatment results and may require further orthodontic treatment. In some cases of atypical growth, the bite may change so much that oral surgery is required to achieve the best possible result.
9. **Residual enamel markings.** Some tooth enamel surfaces are susceptible to being altered by the composite resins used to bond the orthodontic brackets to the teeth. This may result in an altered appearance of the enamel of some of the teeth after the braces are removed.
10. **Headgear.** If improperly handled, headgear may cause injury to the face. Patients are warned not to where the appliance during times of boisterous or competitive activity.
11. **Treatment time.** The total time required to complete treatment may exceed the estimate. Poor co-operation in wearing the appliance or the elastics the required hours per day, poor oral hygiene, broken appliances and missed appointments can lengthen the treatment time and affect the quality of the results.
12. **Co-operation.** Co-operation throughout treatment is your best guarantee of achieving a pleasing smile and a good bite. Failure to co-operate could force the orthodontist to change the procedures and goals of your treatment. As a last resort treatment might have to be suspended. The consequences of early suspension may be worse than no treatment at all.
13. **Complimentary aesthetic dentistry.** If your teeth vary from normal in the number size or shape of teeth, achievement of an ideal result (for example, complete closure of excessive space) may require restorative treatment. The most common types of complementary treatment are cosmetic bonding, crown and bridge restorative dental care and periodontal therapy.
14. **General Health.** General medical problems can affect orthodontic treatment. You should keep your orthodontist informed of any changes to your medical health.

Consent to treatment

I hereby authorised Hunter Valley Orthodontics to carry out orthodontic treatment and any other related procedures deemed necessary including radiographs, for the welfare and treatment of:

Patient's Name

I confirm that the risks attaching to the proposed treatment have been explained to me by my practitioner and that I fully understand those risks and have determined to proceed with treatment.

Signature of patient (or parent/guardian of patients under the age of 18 years)

Signature.....

Name.....

Relationship to Patient.....

Date...../...../.....

Signature of Orthodontist.....